

RESPONSE

Denise Smith PhD, Leanne V. Haller MS, Ron Benedict MS, and Victor Moore PhD

ABSTRACT

was 26.25 beats/min. Exploration and uses revealed that the workload during the training

The impact of resistance training on the cardiovascular system is well documented. The intensity of the training is a key factor in determining the cardiovascular response. The present study examined the cardiovascular response to resistance training in a group of individuals with a history of cardiac rehabilitation. The study was conducted over a period of 12 weeks. The participants were divided into two groups: a control group and an exercise group. The exercise group performed resistance training three times per week. The control group did not exercise. The study found that the exercise group had a significantly lower heart rate during the training compared to the control group. This suggests that resistance training can help to improve cardiovascular fitness in individuals with a history of cardiac rehabilitation.

Key word

cardiovascular response to resistance training

Resistance training is a form of exercise that involves using weights or resistance to build strength and muscle mass. It is a common form of exercise for people who want to improve their physical fitness and health. Resistance training can also help to improve cardiovascular health. The present study examined the cardiovascular response to resistance training in a group of individuals with a history of cardiac rehabilitation. The study was conducted over a period of 12 weeks. The participants were divided into two groups: a control group and an exercise group. The exercise group performed resistance training three times per week. The control group did not exercise. The study found that the exercise group had a significantly lower heart rate during the training compared to the control group. This suggests that resistance training can help to improve cardiovascular fitness in individuals with a history of cardiac rehabilitation.